



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB019Hu01 50 μ g

Recombinant Angiopoietin Like Protein 4 (ANGPTL4)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: His182~Tyr388 (Accession # Q9BY76), with two N-terminal Tags, His-tag and GST-tag.

Host: *E. coli*

Subcellular Location: Secreted, extracellular space, extracellular matrix.

Purity: >95%


Endotoxin Level: <1.0EU per 1 μ g (determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mL Tris, 500mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 6.7

Predicted Molecular Mass:

55.5kDa





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and GST-tag, its sequence is listed below.

MSPILGYWKI KGLVQPTRL L LEYLEEKYEE HLYERDEGDK WRNKKFELGL EFPNLPYYID
GDVKLTQSMA IIRYIADKHN MLGGCPKERA EISMLEGAVL DIRYGVSRIA YSKDFETLKV
DFLSKLP EML KMFEDRLCHK TYLNGDHVTH PDFMLYDALD VVLYMDPMCL DAFPKLVCFK
KRIEAIPIQID KYLKSSKYIA WPLQGQATF GGGDHPPKSD GSTSGSGHHH HHHSAGLVPR
GSTAIGMKET AA AKFERQHM DSPDLGTLEV LFGQPLGSEF- HRLPRDCQE LFQVGERQSG
L F E I Q P Q G S P P F L V N C K M T S D G G W T V I Q R R H D G S V D F N R P W E A Y K A G F G D
PHGEFWLGLE KVHSITGDRN SRLAVQLRDW DGNAELLQFS VHLGGEDTAYSLQLTAPVAG
Q L G A T T V P P S G L S V P F S T W D Q D H D L R R D K N C A K S L S G G W W F G T C S H S N L N
GQYFRSIPQQ RQKLKKGIFW KTRGRYY

[REFERENCES]



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

1. Kim I., *et al.* (2000) *Biochem. J.* 346:603-610.
2. Yoon J.C., *et al.* (2000) *Mol. Cell. Biol.* 20:5343-5349.
3. Ito Y., *et al.* (2003) *Cancer Res.* 63:6651-6657.
4. Clark H.F., *et al.* (2003) *Genome Res.* 13:2265-2270.