

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA050Po01 10µg

Recombinant Insulin Like Growth Factor 1 (IGF1)

Organism Species: Sus scrofa; Porcine (Pig)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Gly49~Ala118

Tags: N-terminal GST-Tag

Accession: P16545

Host: E. coli

Subcellular Location: Secreted.

Purity: >90%

Endotoxin Level: <1.0EU per 1 µ g (determined by the LAL

method).

Formulation: Supplied as lyophilized form in 10ml

PBS, pH7.4, containing 1mM DTT, 5% trehalose, 0.01%

sarcosyl and preservative.


Predicted isoelectric point: 5.9

Predicted Molecular Mass: 34.2kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]





Reconstitute in sterile ddH2O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below. GP ET LC G AE LVD ALQ F VC G D R G FY FN KP T G YG SSS R R AP Q TG IV DE C C FR S C DLRRLEMYCA PLKPAKSA

[REFERENCES]

- 1. Mueller M., Brem G. (1990) Nucleic Acids Res. 18:364-364.
- 2. Tavakkol A., et al. (1988) Mol. Endocrinol. 2:674-681.
- 3. Weller P.A., et al. (1993) J. Mol. Endocrinol. 11:201-211.
- 4. Gregoraszczuk E. L., et al. (2007) J. Reprod. Dev. 53:289-295.