



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YB91571Ra01

Thymus Activation Regulated Chemokine (TARC)

Organism: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

[DESCRIPTION]

Protein Names: Thymus Activation Regulated

Rat TARC

Chemokine

Synonyms: TARC, Ccl17

Species: Rat

Size: 100 μ g

Source: *Escherichia coli*-derived

[PROPERTIES]


Residues: Thr27~Leu93 (Accession # Q9ERE0), with N-terminal His-Tag.

Grade & Purity: >95%, 11kDa as determined by SDS-PAGE reducing conditions.

Formulation: Supplied as lyophilized form in PBS, pH 7.4 containing 5% sucrose, 0.01% sarcosyl.

Endotoxin Level: <1.0 EU per 1 μ g (determined by the LAL method).

Applications: SDS-PAGE; WB; ELISA; IP.





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

(May be suitable for use in other assays to be determined by the end user.)

Predicted Molecular Mass: 9.3kDa

Predicted isoelectric point: 9.5

[PREPARATION]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MGHHHHHSGSEF- TNVG RECCLDYFKG AIPIRKLVTV FRTSVECPKD AIVFETVQGR
LICTDPKDKH VKKAIRHLKN QRL