

YBF656Hu01 100µg

Recombinant Phosphodiesterase 10A (PDE10A)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

11th Edition (Revised in May, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: E. coli

Residues: Asn508~Asp779 Tags: N-terminal His-Tag

Tissue Specificity: Brain, Testis. Subcellular Location: Cytoplasm.

Purity: >96%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; Purification; Amine

Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.0

Predicted Molecular Mass: 34.7kDa

Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8_oC for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37_°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

NYR RVPYHNWKHA VTVAHCMYAI LONNHTLFTD LERKGLLIAC LCHDLDHRGF SNSYLQKFDH PLAALYSTST MEQHHFSQTV SILQLEGHNI FSTLSSSEYE QVLEIIRKAI IATDLALYFG NRKQLEEMYQ TGSLNLNNQS HRDRVIGLMM TACDLCSVTK LWPVTKLTAN DIYAEFWAEG DEMKKLGIOP IPMMDRDKKD EVPQGQLGFY NAVAIPCYTT LTQILPPTEP LLKACRDNLS QWEKVIRGEE TATWISSPSV AQKAAASED

[IDENTIFICATION]




Figure 1. SDS-PAGE



EL:4006-871-227 Web:www.ybi o.net Email:shy bio@126.com