



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA225Mu01 50 μ g

Atrial Natriuretic Peptide (ANP)

Organism Species: *Mus musculus* (Mouse)

*Instruction
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul,
2013)

[PROPERTIES]

Residues: Asn25~Arg152 (Accession # P05125), with two N-terminal Tags, His-tag and GST-tag.

Host: *E. coli*


Subcellular Location: Secreted.

Purity: >95%

Endotoxin Level: <1.0EU per 1 μ g (determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4 containing 5% sucrose, 0.01% sarcosyl.

Predicted isoelectric point: 6.1 Predicted



The possible reasons that the actual band size differs from the predicted are as follows:

Molecular Mass: 46.0kDa

Accurate Molecular Mass: 44kDa as determined by SDS-PAGE reducing conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note:



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and GST-tag, its sequence is listed below.

MSPILGYWKI KGLVQPTRLL LEYLEEKYEE HLYERDEGDK WRNKKFELGL EFPNLPYYID
GDVKLTQSM A IIRYIADKHN MLGGCPKERA EISMLEGAVL DIRYGVSRIA YSKDFETLKV
DFLSKLP EML KMFEDRLCHK TYLNGDHVTH PDFMLYDALD VVLYMDPMCL DAFPKLVCFK
KRIEAIPQID KYLKSSKYIA WPLQGQWATF GGGDHPPKSD GSTSGSGHHH HHHSAGLVPR
GSTAIGMKET AAKFERQHM DSPDLGTLEV LFQGPLGSEF-NPVYSA VSNTDLMDFK



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

NLLDHLEEKM PVEDEVMPQ ALSEQTEEAG AALSSLPEVP PWTGEVNPPL RDGSALGRSP

WD P S D R S A LL KS K L R A L L A G PR S LR R S SC F GG R I D R I G A Q SG LG C N S F RY RR

[REFERENCES]

1. Cui Y., *et al.* (2012) Nature 484:246-250.
2. Chan J.C., *et al.* (2005) Proc. Natl. Acad. Sci. U.S.A. 102:785-790.
3. Madhani M., *et al.* (2003) Br. J. Pharmacol. 139:1289-1296.
4. Carninci P., *et al.* (2005) Science 309:1559-1563.