



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBD111Ra01 50 μ g

Recombinant Gastric Intrinsic Factor (GIF)

Organism Species: *Rattus norvegicus* (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Arg196~Phe414

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: P17267

Host: *E. coli*

Subcellular Location: Secreted.

Purity: >95%

Endotoxin Level: <1.0EU per 1 μ g
(determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mL


Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM
DTT, 0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 6.6

Predicted Molecular Mass: 29.5kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)





[USAGE]

Reconstitute in ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

RI PV G SQ EN Y RD LF G QA LK VI VD NI SLR IK AD G II GD IY ST G LAM QA LS VT PE QP
TKEWDCEKTM YTILKEIKQG KFNPMISIAQ ILPSLKGKTY LDVPQVTCGP DHEVPPTLTD
YPTPVPTSIS NITVIYTINN QLRGVDLLFN VTIEVSVKSG SVLLAVLEEA QRRNHMFKFE
TT M T S W G L I V S S I N N I A E N V K H K T Y W E F L S G K T P L G E G V A Y Y I P F N Y E H I T A N F

[REFERENCES]

1. Dieckgraefe B.K., *et al.* (1988) Proc. Natl. Acad. Sci. U.S.A. 85:46-50.
2. Maeda M., *et al.* (1995) J. Biochem. 117:1305-1311.
3. BOASS A., WILSON T.H. (1964) Am. J. Physiol. 206:783-786.
4. Khadapkar S.V., Vakil U.K. (1982) Biochim. Biophys. Acta 719:619-632.